

Marginal Pinning of Quenched Random Polymers

D.A. Gorokhov and G. Blatter

Theoretische Physik, ETH-Hönggerberg, CH-8093 Zürich, Switzerland
(February 1, 2008)

An elastic string embedded in 3D space and subject to a short-range correlated random potential exhibits marginal pinning at high temperatures, with the pinning length $L_c(T)$ becoming exponentially sensitive to temperature. Using a functional renormalization group (FRG) approach we find $L_c(T) \propto \exp[(32/\pi)(T/T_{dp})^3]$, with T_{dp} the depinning temperature. A slow decay of disorder correlations as it appears in the problem of flux line pinning in superconductors modifies this result, $\ln L_c(T) \propto T^{3/2}$.

Elastic manifolds subject to a disorder potential [1] define a rich and challenging problem in modern classical statistical mechanics, with numerous applications in condensed matter physics [1,2,3,4]. Fundamental questions arise regarding the relevance of the disorder [5] and its quantitative effects on the manifold's quenched fluctuations on short [5] and large scales [1]. Such static results then determine, via scaling arguments, the creep type motion of the elastic manifolds under the action of an external force, rendering these studies also relevant in the context of applied physics, e.g., plastic deformations in metals [3] or dissipation in superconductors [2]. Classifying the problems through the dimensionality d of the manifold and the number n of transverse dimensions, the $(1+n)$ -dimensional problem describing strings moving in n directions, also known as the random polymer problem, is particularly interesting [6,7]. Here, depending on the number of transverse dimensions n , the temperature induced fluctuations compete in various ways with the fluctuations due to quenched disorder. In this Letter, we analyze the competition between elasticity, disorder, and temperature for the $(1+n)$ -string problem. In particular, we present a functional renormalization group analysis of the $(1+2)$ -dimensional problem (a string in 3D space), where the disorder turns marginal at high temperatures; we determine the depinning temperature T_{dp} above which thermal smoothing leads to a collapse of the effective disorder strength and the temperature dependent pinning length $L_c(T)$ beyond which the disorder dominates over the elasticity.

The free energy $F(\mathbf{u}, L) = -T \ln Z(\mathbf{u}, L)$ of a string starting and ending at the points $(\mathbf{0}, 0)$ and $(\mathbf{u}, L)$ [the first (second) coordinate denotes the displacement of (position along) the string]] is given by the partition function

$$Z = \int_{(\mathbf{0},0)}^{(\mathbf{u},L)} \mathcal{D}[\mathbf{u}'] \exp \left\{ -\frac{1}{T} \int_0^L dz' \left[\frac{\epsilon}{2} \left(\frac{\partial \mathbf{u}'}{\partial z'} \right)^2 + U(\mathbf{u}', z') \right] \right\},$$

with ϵ the elasticity of the string. The disorder potential $U(\mathbf{u}, z)$ is assumed to be a random gaussian variable with zero mean and a correlator

$$\langle U(\mathbf{u}, z) U(\mathbf{u}', z') \rangle = K_0(|\mathbf{u} - \mathbf{u}'|) \delta(z - z'), \quad (1)$$

with $K_0(u)$ a smooth function decaying on the scale ξ . The partition function Z describes the competition between the elastic energy $(\epsilon/2)(\partial_z \mathbf{u})^2 > 0$ and the disorder potential $U(\mathbf{u}, z)$, from which the string can gain energy by choosing minima with $U(\mathbf{u}, z) < 0$.




FIG. 1. Schematic picture of the FRG flow for the marginal case with $n = 2$: At temperatures below depinning ($T < T_{dp} \approx (\epsilon K_0(0)\xi^2)^{1/3}$) the flow proceeds through the Larkin regime (with a fixed point characterized by the wandering exponent $\zeta_L = 3/2$) towards the random manifold fixed point with its non-trivial exponent $\zeta_2 \approx 5/8$. At high temperatures $T > T_{dp}$ the flow starts out in the thermal region with $\zeta_{th} = 1/2$. The linear flow towards the ‘Boltzmann’ fixed point (BFP) rapidly erases the details in the initial condition and produces a gaussian correlator $K_l(u) \sim g_l \exp(-u^2 \epsilon / 4\pi \xi T)$ with weight g_l . The non-linear terms in the FRG produce an overall growth of the correlator while the linear terms keep its shape to the ‘Boltzmann’ form (adiabatic pumping). The sharp increase of the prefactor g_l at l_c defines the pinning length $L_c(T)$, beyond which the flow enters the random manifold regime with a rapid change of the thermal exponent to the random manifold exponent ($\zeta_{th} = 1/2 \rightarrow \zeta(l) \rightarrow \zeta_2 \approx 5/8$) and a renormalization of the temperature towards zero. The flow ends in the zero temperature random manifold fixed point.

The quantity characterizing the behavior of the string is the displacement correlator $\langle u^2(L) \rangle \equiv \langle [\mathbf{u}(L) - \mathbf{u}(0)]^2 \rangle \propto L^{2\zeta}$ describing its wandering with distance L . We distinguish between the perturbative (Larkin [5]) regime at short scales $L < L_c$ and the random manifold regime [8] at $L > L_c$, as well as temperatures below and above the depinning temperature $T_{dp} \approx (\epsilon K_0(0)\xi^2)^{1/3}$, see Fig. 1: For $T < T_{dp}$ the wandering exponent $\zeta_L = 3/2$ at

small distances $L < L_c$ (see [2]), while at large distances $L > L_c$, ζ_n depends on the dimensionality of the space: $\zeta_1 = 2/3$ is an exact result [7] while for $n = 2$ numerical simulations [9] give a value $\zeta_2 \approx 5/8$. The crossover length L_c separating these two regimes has the form [2] $L_c \approx [\epsilon^2 \xi^4 / K_0(0)]^{1/3}$. At high temperatures $T > T_{dp}$ the short scale behavior is subject to thermal smoothing, resulting in a thermal wandering with $\zeta_{th} = 1/2$, while at large scales $L > L_c(T)$ the disorder induced line wandering characterized by ζ_n prevails. The crossover length $L_c(T)$ increases algebraically with temperature $L_c(T) \sim L_c(T/T_{dp})^5$ for the $(1+1)$ -case [3], while for the $(1+2)$ -dimensional problem $L_c(T)$ is exponentially sensitive to temperature with [10] $L_c(T) \sim A(T) \exp[C(T/T_{dp})^3]$, and C a constant of order unity. It turns out that the usual perturbative methods neither provide the value of C nor that of the prefactor $A(T)$. The exponential sensitivity of the pinning length $L_c(T)$ to the temperature T is related to the appearance of a phase transition in dimensions $n > 2$, separating a low temperature disorder dominated phase from a high temperature thermal phase characterized by ζ_{th} [11]. The $(1+2)$ -problem then corresponds to the lower critical dimension for this phase transition and thus exhibits *marginal* behavior.

The main goal of the present work is to develop a consistent scheme for calculating the pinning length $L_c(T)$ in the $(1+2)$ -problem specifying the coefficient C in the exponent. While this goal can be achieved within a one-loop calculation including three-replica terms, it turns out that the determination of the prefactor $A(T)$ requires a two-loop analysis (accounting for four-replica terms) which is beyond the scope of this letter. We first discuss perturbative methods for calculating $L_c(T)$ at high temperatures and different dimensions n . These techniques fail for the $(1+2)$ -problem and we will then turn to the more powerful renormalization group method.

Perturbation Theory: With the weak disorder providing the small parameter $\Delta = \int d^n u K_0(u)$ we expand the k -fold replicated Green function (see [12]) $G(\mathbf{u}_1, \dots, \mathbf{u}_k; L) \equiv \langle Z(\mathbf{u}_1, L) \dots Z(\mathbf{u}_k, L) \rangle$ and calculate the mean squared displacement $\langle u^2(L) \rangle = \lim_{k \rightarrow 0} \int d^n u d^n u_2 \dots d^n u_k u^2 G(\mathbf{u}, \mathbf{u}_2, \dots, \mathbf{u}_k; L)$. The disorder is relevant when $\langle u^2(L \rightarrow \infty) \rangle$ grows faster than the thermal wandering $\langle u^2(L) \rangle_{th} = (nT/\epsilon)L$ which is dominant at small lengths; the comparison of the two expressions then defines the characteristic crossover length $L_c(T)$, $(nT/\epsilon)L_c(T) \sim \langle u^2(L_c(T)) \rangle$. To lowest order the mean squared displacement takes the form

$$\langle u^2(L) \rangle = \frac{nTL}{\epsilon} \left[1 + \frac{(\Delta/T^2)(\epsilon/T)^{n/2}}{(4\pi)^{n/2}(4-n)} L^{1-n/2} \right]; \quad (2)$$

for $n < 2$ the disorder corrections grow more rapidly than the thermal wandering and we obtain [3] $L_c(T) \sim L_c [T/(\epsilon \Delta \xi)^{1/3}]^5$ for $n = 1$. For $2 < n < 4$ the contribution from the random potential goes to zero at large

scales and the disorder is irrelevant [11]. The case $n = 2$ is marginal: to lowest order the disorder correction is L -independent and the next order term $\propto \Delta^2$ provides a logarithmic correction,

$$\langle u^2(L) \rangle = \frac{2TL}{\epsilon} \left[1 + \frac{1}{8\pi} \frac{\epsilon \Delta}{T^3} + \frac{1}{16\pi^2} \frac{\epsilon^2 \Delta^2}{T^6} \ln(L/\xi) \right], \quad (3)$$

producing an exponential temperature dependence $L_c(T) \propto \exp[CT^3/\epsilon \Delta]$ with C of order unity (we compare terms $\propto \Delta^2$ and $\propto \Delta^3$). The arbitrariness of the criterion prevents us from finding the coefficient in the exponent and the prefactor and we have to develop a more systematic way in order to deal with the problem.

Renormalization group: The basic idea is to construct the renormalized effective correlator K_l of the disorder potential describing the behavior of the manifold on short and large scales. With the disorder becoming irrelevant for (internal) dimensionalities $d > 4$ [5], the effect of disorder is analyzed within an $\epsilon = 4 - d$ -expansion. Starting from short scales, the RG flow goes through a special point where K_l becomes singular. This point is identified with the pinning length L_c of the string [13]. Its physical meaning is clear: the fact that the correlator becomes singular at L_c implies that the perturbation theory breaks down and the manifold splits up into independently pinned domains of size L_c . By way of introduction we briefly derive the zero-temperature pinning length L_c , starting from the one-loop FRG equation [8,14]

$$\partial_l K_l = (4 - d - 4\zeta - n\zeta)K_l + \zeta \nabla \cdot (\mathbf{u} K_l) + I[K_l^{\mu\nu}(\mathbf{u})K_l^{\mu\nu}(\mathbf{u})/2 - K_l^{\mu\nu}(\mathbf{u})K_l^{\mu\nu}(\mathbf{0})], \quad (4)$$

with $\zeta(l) = (1/2)\partial \ln \langle u^2 \rangle / \partial \ln L$ the wandering exponent, $I = A_d/(2\pi)^d \epsilon^2 \Lambda^{4-d}$ ($A_d = 2\pi^{d/2}/\Gamma(d/2)$, and $\Lambda^{-1} \sim \xi$ is the short scale cutoff), and the indices μ and ν denote derivatives with respect to the Cartesian coordinates u_μ and u_ν . The RG variable l is related to the length L via $l = \ln \Lambda L$. We proceed with differentiating (4) four times with respect to u_κ . Substituting $u = 0$, the wandering exponent ζ drops out and we obtain

$$\partial_l \Gamma_l = (4 - d)\Gamma_l + [I(n + 8)/3]\Gamma_l^2, \quad (5)$$

where $\Gamma_l = \partial^4 K_l / \partial u_\kappa^4|_{\mathbf{u}=0}$. Integrating (5) we encounter a divergence at $l_c \approx \ln[3(4 - d)/I(n + 8)\Gamma_0]/(4 - d)$ (we assume weak disorder with $I\Gamma_0 \ll 1$), allowing us to define the collective pinning or Larkin length $L_c = \Lambda^{-1} \exp(l_c) \approx [\epsilon^2 \xi^4 / K(0)]^{1/(4-d)}$ where simple perturbation theory breaks down. The FRG equation (4) allows us to further characterize the Larkin regime: On short scales we can neglect the non-linear term and an expansion of the correlator $K_l(u) \sim \alpha(l)u^2/2$ produces the wandering exponent $\zeta_L = (4 - d)/2$ at the Larkin fixed point [15]. Note that, although Eq. (4) is written in the form of an ϵ -expansion, the result for L_c is valid for any ϵ as long as we investigate the short scale behavior, where the functional renormalization group is just

a way to sum the perturbation expansion. Indeed, one can show that in the non-marginal situation higher order loop corrections to Eq. (4) do not change the result for L_c , while in the marginal case (e.g., for $d = 4$ at $T = 0$) we have to account for two-loop corrections.

Next, we proceed to study finite temperatures, where besides a simple diffusive term $\propto T\Delta K_l$, new three- and higher replica terms are generated describing non-gaussian fluctuations in the disorder potential U . Generalizing the disorder statistics to include non-gaussian terms and replicating n times we arrive at a new Hamiltonian $\mathcal{H}_n = \mathcal{H}_{\text{free}} + \mathcal{H}_{\text{int}}$ with $\mathcal{H}_{\text{free}} = \int d^d z \sum_{\alpha} (\epsilon/2)(\partial_{\mathbf{z}} \mathbf{u}_{\alpha})^2$ and

$$\frac{\mathcal{H}_{\text{int}}}{T} = - \int d^d z \left[\sum_{\alpha, \beta} \frac{K(\mathbf{u}_{\alpha\beta})}{2T^2} + \sum_{\alpha, \beta, \gamma} \frac{S(\mathbf{u}_{\alpha\beta}, \mathbf{u}_{\alpha\gamma})}{T^3} + \dots \right].$$

Here, $\mathbf{u}_{\alpha\beta} \equiv \mathbf{u}_{\alpha} - \mathbf{u}_{\beta}$ and $S(\mathbf{u}, \bar{\mathbf{u}})$ is the three-replica term describing non-gaussian fluctuations in the disorder potential (the $\dots$ stand for higher-replica terms depending on three or more variables). At finite temperatures, a consistent analysis to one loop order requires to include both two- and three-replica terms $K(\mathbf{u})$ and $S(\mathbf{u}, \bar{\mathbf{u}})$ in the FRG flow. After a second order cumulant expansion in the two- and three-body interaction $\mathcal{H}_{\text{int}}$, we proceed with the standard momentum shell RG, expand the $\mathbf{u}$ -fields into Fourier series and perform the integration over fast modes. The resulting equations take the form

$$\begin{aligned} \partial_l K_l(\mathbf{u}) = & (4-d-4\zeta-n\zeta)K_l + \zeta \nabla \cdot (\mathbf{u} K_l) + I' T K_l^{\mu\mu} \\ & + I[(1/2)K_l^{\mu\nu}(\mathbf{u})K_l^{\mu\nu}(\mathbf{u})/2 - K_l^{\mu\nu}(\mathbf{u})K_l^{\mu\nu}(\mathbf{0})] \\ & - 2I'T[S_l^{\mu\mu}(\mathbf{0}, \mathbf{u}) + S_l^{\bar{\mu}\bar{\mu}}(\mathbf{u}, \mathbf{0}) - [\partial_{\mathbf{u}} \cdot \partial_{\bar{\mathbf{u}}} S_l](\mathbf{u}, \mathbf{u}) \\ & + [\partial_{\mathbf{u}} \cdot \partial_{\bar{\mathbf{u}}} S_l](\mathbf{0}, \mathbf{u}) + [\partial_{\mathbf{u}} \cdot \partial_{\bar{\mathbf{u}}} S_l](\mathbf{u}, \mathbf{0})], \end{aligned} \quad (6)$$

$$\begin{aligned} \partial_l S_l(\mathbf{u}, \bar{\mathbf{u}}) = & (6-2d-6\zeta-2n\zeta)S_l + \zeta[\nabla \cdot (\mathbf{u} S_l) + \bar{\nabla} \cdot (\bar{\mathbf{u}} S_l)] \\ & + I'T[S_l^{\mu\mu} + S_l^{\bar{\mu}\bar{\mu}} + \partial_{\mathbf{u}} \cdot \partial_{\bar{\mathbf{u}}} S_l] \\ & + I T K_l^{\mu\nu}(\mathbf{u}) K_l^{\mu\nu}(\bar{\mathbf{u}})/4, \end{aligned} \quad (7)$$

$$\partial_l T_l = (2-d-2\zeta)T_l, \quad (8)$$

where $I' = I\epsilon\Lambda^2$ and the superscripts μ ($\bar{\mu}$) denote derivatives with respect to the first (second) variable in $S_l(\mathbf{u}, \bar{\mathbf{u}})$ (implicit summation over double indices is assumed); similarly, the gradients ∇ ($\bar{\nabla}$) denote derivatives with respect to $\mathbf{u}$ ($\bar{\mathbf{u}}$). The system (6) – (8) has to be solved with initial conditions $K_{l=0}(\mathbf{u}) = K_0(\mathbf{u})$, $S_{l=0}(\mathbf{u}, \bar{\mathbf{u}}) = 0$ (the bare potential is gaussian), and $T_{l=0} = T$. Note that the three-replica term is driven by a source $\propto T K_l^2$ and hence is irrelevant in the $T = 0$ analysis. Furthermore, four- and higher replica terms can be omitted within the present one-loop analysis: with $K_l(\mathbf{u}) = \mathcal{O}(\Delta)$ the three-replica term $S_l(\mathbf{u}, \bar{\mathbf{u}}) = \mathcal{O}(\Delta^2)$, see (7), while the four-replica term is driven by a term $\propto K_l^3$ and hence $D(\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3) = \mathcal{O}(\Delta^3)$. However, going to second-loop order the four-replica term D_l has to be included.

The above system (6) – (8) can be solved asymptotically exactly for weak disorder and we proceed with the

analysis for the elastic string, $d = 1$. On short scales the behavior of the string is dominated by thermal fluctuations and the wandering exponent is equal to $\zeta_{\text{th}} = 1/2$; as a result, the temperature is not renormalized, see Eq. (8). On the other hand, on large scales ζ approaches the value $\zeta_2 \approx 5/8$ and the behavior of the string is governed by the zero temperature fixed point. It turns out that the temperature starts renormalizing only close to $L_c(T)$, allowing us to fix $\zeta = 1/2$ in Eqs. (6) – (8). The non-linear and three-replica terms are of second order in the disorder strength and thus unimportant at small scales $l \ll l_c$, allowing us to neglect the terms $\propto K_l^2$ and $\propto S_l$ in (6) in the initial stage of the flow. The Ansatz

$$K_l(u) = P_l(u) \exp[(1-n/2)l], \quad (9)$$

then maps (6) to the Fokker-Planck equation describing the ‘time’ evolution of the probability distribution function P_l for an overdamped particle in the harmonic potential $V(u) = u^2/4$ at a ‘temperature’ $T' = T/\pi\epsilon\Lambda$, $\partial_l P_l = \nabla \cdot (\mathbf{u} P_l)/2 + T' \Delta P_l$. At high temperatures T , the distribution function $P_l(u)$ rapidly approaches equilibrium and takes the form of a Boltzmann distribution,

$$P_B(u) = \Delta(4\pi T')^{-n/2} \exp(-u^2/4T'). \quad (10)$$

The function $P_B(u)$ depends on the disorder potential only via the integral weight $\Delta = \int d^n u K_0(u)$, i.e., the evolution erases the details of the initial condition. Depending on the dimensionality n , the following situations arise (c.f., (9)): For $n > 2$ the solution of (6) decreases with increasing l and the disorder renormalizes to zero, in agreement with the existence of a high temperature phase when $n > 2$, see [11]. For $n < 2$ the correlator $K_l(u) = \exp[(1-n/2)l]P_B(u)$ grows (until reaching the size of the non-linear term) and the disorder is relevant. Comparing terms linear and quadratic in K_l we find the collective pinning length $L_c(T) \simeq L_c[T/(\epsilon\Delta\xi)^{1/3}]^5$. Most interesting is the marginal case $n = 2$ where we find a ‘Boltzmann’ fixed point. In order to obtain $L_c(T)$ we need to take into account the terms $\mathcal{O}(\Delta^2)$ effectively pumping additional ‘particles’ into the system and leading to an overall growth in the disorder strength $\Delta_l = \int d^n u K_l(u)$. However, for small Δ_l the strong diffusion will relax this ‘pumping’ towards the ‘Boltzmann’ distribution, i.e., the solution of Eq. (6) keeps the form

$$K_l(\mathbf{u}) = g_l (T'^2/I) \exp(-\mathbf{u}^2/4T'), \quad (11)$$

with the dimensionless coupling $g_l = (\pi/4)\Delta_l\epsilon/T^3$. After diagonalizing the term $[S_l^{\mu\mu} + S_l^{\bar{\mu}\bar{\mu}} + \partial_{\mathbf{u}} \cdot \partial_{\bar{\mathbf{u}}} S_l]$ in (7) we find its Green function $G_S(\mathbf{u}, \bar{\mathbf{u}}; \mathbf{u}', \bar{\mathbf{u}}'; l) = 1/12[\pi I'T(1-e^{-l})]^2 \exp\{-[(\mathbf{u}-\mathbf{u}'e^{-l/2})^2 + (\bar{\mathbf{u}}-\bar{\mathbf{u}}'e^{-l/2})^2 + (\mathbf{u}-\mathbf{u}'e^{-l/2}) \cdot (\bar{\mathbf{u}}-\bar{\mathbf{u}}'e^{-l/2})]/3I'T(1-e^{-l})\}$ and obtain the three-replica term $S_l(\mathbf{u}, \bar{\mathbf{u}})$ by simple integration. Substituting S_l back into (6) and integrating over $\mathbf{u}$ we find that the three-replica terms mutually cancel and thus do not contribute

to the flow of g_l . The amplitude g_l then is determined by the remaining non-linear terms alone,

$$\partial_l g_l = g_l^2/8, \quad (12)$$

with the solution $g_l = g_0/(1 - g_0 l/8)$ as derived from the initial condition $g_0 = (\pi/4)(\epsilon\Delta/T^3)$. At $l_c = 8/g_0$ the amplitude $g_{l_c} \approx 1$ and we find the pinning length

$$L_c(T) \sim \exp[(32/\pi)(T^3/T_{\text{dp}}^3)], \quad T_{\text{dp}}^3 = \epsilon\Delta, \quad (13)$$

the main result of the paper [16]: the one-loop FRG analysis allows us to fix the coefficient C in front of the parameter $1/g_0$ in the exponent. Although the criterion $g_{l_c} \simeq 1$ might seem arbitrary it is sufficient to find the correct asymptotic value of the RG variable l_c as long as the initial value g_0 is small. Indeed, the solution g_l remains small in most of the interval $[0, l_c]$ and grows rapidly only very close to l_c .

An interesting alternative is offered by mapping the random polymer problem to the Kardar-Parisi-Zhang equation [17], where the dynamic renormalization group (DRG) [18] provides excellent results for the $n = 1$ problem [19]. The weight $\tilde{g}$ in the (white noise) Ansatz $K_{\text{wn}} = \tilde{g}\delta^2(\mathbf{u})$ replacing (11) obeys the equation [20] $\partial_l \tilde{g} = (2 - n)\tilde{g} + [2(2n - 3)/n]\tilde{g}^2$, with $\tilde{g}(0) = (A_n/(2\pi)^n)\epsilon\Delta/T^3$ (the prime indicates scaling along the transverse dimension). For $n = 2$ the solution ‘explodes’ at $l'_c = 2\pi(T/T_{\text{dp}})^3$, implying that $L_c(T) \sim \exp[4\pi(T/T_{\text{dp}})^3]$ [21], different from Eq. (13). Note that, in contrast to the FRG, the DRG scheme does not renormalize the correlator *function*. Here, we argue in favor of our result (13): While the DRG technique produces accurate results (including a random manifold fixed point) in dimensions $n < 3/2$, the results are less convincing in larger dimensions $n > 3/2$, where the fixed point gives way to a divergence. On the other hand, the FRG technique features both an apparent divergence at l_c and a random manifold fixed point as $l \rightarrow \infty$ in all dimensions $n \leq 2$, allowing for a consistent physical interpretation of the results in terms of the diagram sketched in Fig. 1.

Above we have assumed that the integral $\int d^n u K_0(u)$ converges, however, flux lines in superconductors exhibit a long range interaction with the disorder potential spoiling this assumption: for values u exceeding the coherence length ξ , the correlator obeys the asymptotics $K_0(u) \sim K_0(0)(\xi^2/u^2)\ln(u/\xi)$ and the integral $\int_{u < u^*} d^n u K_0(u)$ diverges as $K_0(0)\xi^2 \ln^2(u^*/\xi)$. We estimate $L_c(T)$ for this case: Neglecting the non-linear term in the first stage of the RG transformation, we approximate the solution of the Fokker-Planck equation in the region $u^2 \lesssim T\xi/\epsilon$ by $[\Delta_l/4\pi T'] \exp[-u^2/4T']$ [c.f., Eq. (10)], with an l -dependent function Δ_l , a consequence of the divergence of the integral $\int d^n u K_0(u)$. The weight Δ_l collects those particles reaching the central region on scale l : a displacement $u^* \gg \sqrt{T\xi/\epsilon}$ of the harmonic oscillator takes a ‘time’ $l^* \sim \ln(u^*/\sqrt{T\xi/\epsilon})$ to reach the region $u \lesssim \sqrt{T\xi/\epsilon}$

and hence $\Delta_{l^*} \sim K_0(0)\xi^2 l^{*2}$. The dimensionless coupling g_l then grows already in the linear regime due to the ‘pumping’ from large distances. The pinning length follows from the condition $g_{l_c} = \epsilon\Delta_{l_c}/T^3 \approx 1$ and we obtain $L_c(T) \sim \exp[C(T^3/\epsilon K_0(0)\xi^2)^{1/2}]$ with C of order unity. The same result follows from a perturbative treatment.

In order to find the prefactor $A(T)$ in the expression (13) for $L_c(T)$ we have to account for $\mathcal{O}(\Delta^3)$ corrections, resulting in a flow $\partial_l g_l = g_l^2/8 + \eta g_l^3$ with $\eta > 0$ a constant. This equation explodes at $l_c = 8/g_0 - 64\eta \ln g_0$, producing a prefactor of the form $A(T) \simeq \Lambda^{-1}(T_{\text{dp}}/T)^{192\eta}$. While the two-loop contribution $\eta_{2-1} = 275/5184$ to η can be found from [13], the part arising from the four-replica term is more difficult to obtain — whether the four-replica terms mutually cancel (e.g., due to some symmetry), as was the case in the one-loop analysis above, remains an interesting and open problem.

We thank L. Balents, D.S. Fisher, E. Frey, V.B. Geshkenbein, T. Giamarchi, Th. Hwa, O. Narayan, V.M. Vinokur, and J. Zinn-Justin for helpful discussions.

-
- [1] T. Halpin-Healy and Y.-C. Zhang, Phys. Rep. **254**, 215 (1995).
 - [2] G. Blatter *et al.*, Rev. Mod. Phys. **66**, 1125 (1994).
 - [3] L.B. Ioffe and V.M. Vinokur, J. Phys. C **20**, 6149 (1987).
 - [4] G. Grüner, Rev. Mod. Phys. **60**, 1129 (1988).
 - [5] A. I. Larkin, Zh. Eksp. Teor. Fiz. **58**, 1466 (1970) [Sov. Phys. JETP **31**, 784 (1970)].
 - [6] D.A. Huse and C.L. Henley, Phys. Rev. Lett. **54**, 2708 (1985).
 - [7] D.A. Huse *et al.*, Phys. Rev. Lett. **55**, 2924 (1985).
 - [8] D.S. Fisher, Phys. Rev. Lett. **56**, 1964 (1986).
 - [9] T. Ala-Nissila and O. Venäläinen, J. Stat. Phys. **76**, 1083 (1994).
 - [10] M.V. Feigel'man and V.M. Vinokur, Phys. Rev. B **41**, 8986 (1990).
 - [11] J.Z. Imbrie and T. Spencer, J. Stat. Phys. **52**, 609 (1988).
 - [12] D.A. Gorokhov and G. Blatter, Phys. Rev. Lett. **82**, 2705 (1999).
 - [13] H. Bucheli *et al.*, Phys. Rev. B **57**, 7642 (1998); D.A. Gorokhov and G. Blatter, Phys. Rev. B **59**, 32 (1999).
 - [14] L. Balents and D.S. Fisher, Phys. Rev. B **48**, 5949 (1993).
 - [15] T. Giamarchi and P. LeDoussal, Phys. Rev. B **52**, 1242 (1995).
 - [16] Previous results $L_c(T) \propto T^4$ for $n = 1$ and $\ln L_c(T) \propto T^2$ for $n = 2$ found by Kim *et al.* [Phys. Rev. A **44**, 4782 (1991)] disagree with those from perturbation theory. Furthermore, the result $L_c(T) \propto \exp[16\pi(T/T_{\text{dp}})^3]$ by Lebedev [Europhys. Lett. **16**, 1 (1991)] is based on the summation of the main terms in the perturbation series, which, however, have been chosen on the basis of a simplified rule.
 - [17] M. Kardar *et al.*, Phys. Rev. Lett. **56**, 889 (1986).
 - [18] D. Forster *et al.*, Phys. Rev. A **16**, 732 (1977).
 - [19] E. Medina *et al.*, Phys. Rev. A **39**, 3053 (1989).
 - [20] E. Frey and U.C. Tauber, Phys. Rev. E **50**, 1024 (1994).
 - [21] D. Fisher and D.A. Huse, Phys. Rev. B **43**, 10728 (1991).